

UNCLASSIFIED

AD NUMBER	
AD144275	
CLASSIFICATION CHANGES	
TO:	UNCLASSIFIED
FROM:	SECRET
LIMITATION CHANGES	
TO: Approved for public release; distribution is unlimited. Document partially illegible.	
FROM: Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; FEB 1956. Other requests shall be referred to Deputy Chief of Staff Research and Development, Department of the Air Force, Washington, DC 20330.	
AUTHORITY	
13 Nov 1961 per ltr; Rand ltr, 21 Feb 1967	

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED

AD

144275

CLASSIFICATION CHANGED
TO: **UNCLASSIFIED**
FROM: **SECRET**
AUTHORITY:

USAF 21 Dec. 60
Per letter 13 Nov. 61



UNCLASSIFIED

FORMERLY RESTRICTED DATA

as Restricted Data in foreign dissemination.
Section 144b, Atomic Energy Act, 1954

SECRET

RM-1583-1
2-3-56
-5-

They also represent burst in a homogeneous sea-level atmosphere with standard ambient pressure $P_0 = 14.7$ psi, density $\rho_0 = .00129$ gm/cc, and temperature $T_0 = 273.2$ °Kelvin.

Any other yield or ambient conditions are achieved by multiplying all distances by $(W/\pi)^{1/3}$ and all times by $(W/\pi)^{1/3} \tau^{-1/2}$ where W is the yield in megatons, π is the ratio of ambient pressure (P_a) to standard pressure (P_0), and τ is the ratio of ambient temperature (T_a) to standard temperature (T_0). Under this scaling the various pressure ratios will be the same, but pressures must be multiplied by π , velocities (Mach numbers) by $\tau^{1/2}$ and temperatures by τ .

Figure 1. is a plot of peak overpressure in atmospheres (14.7 psi) versus range in kilofeet.

Figure 2. is a similar plot of the peak dynamic pressure ($1/2 \rho u^2$) in atmospheres versus range in kilofeet. It should be noted that the usual Hugoniot relation for an ideal gas of $\gamma = 1.4$, viz. $Q_s = \frac{5}{2} \frac{\Delta P_s^2}{7P_0 + \Delta P_s}$ is valid only for small pressures ($\Delta P_s < 3$ atmos), and that for higher pressures the peak dynamic pressures are greater than this (characteristic of a smaller γ).

Figure 3. is a plot of particle velocity (u_s) and shock velocity (U_s) in units of ambient sonic velocity, i.e., a plot of particle and shock Mach numbers. Also plotted is the shock density (ρ_s) in units of ambient density (ρ_0) versus range in kilofeet.

Figures 4 - 21 are plots of overpressure (ΔP) in atmos, dynamic pressure (Q) in atmos, particle Mach number (U), density ratio (ρ), and temperature ratio

SECRET

S E C R E T

UNCLASSIFIED

Work of RAND Corp. is in the Public Domain
Under Patent Rights Act, 1951

U.S. AIR FORCE

Project RAND

RESEARCH MEMORANDUM



This report contains information affecting the national defense of the United States within the meaning of the espionage laws, Title 18, U.S.C., Sec. 793 and 794, and the transmission or the revelation of its contents in any manner to an unauthorized person is prohibited by law.

THIS PAGE FOR INFORMATION PURPOSES

This is a working paper, because it may be amended, revised, or otherwise changed after issuance. It is not to be used as a basis of action. The views, conclusions, and recommendations expressed herein are those of the author and do not represent the official position of the RAND Corporation.

RAND

Corporation

S E C R E T

571A

46653

SECRET
FORMERLY RESTRICTED DATA

Handle As Restricted Data In Foreign Dissemination
Section 144b, Atomic Energy Act, 1954

U. S. AIR FORCE
PROJECT RAND
RESEARCH MEMORANDUM

CLOSE-IN H-BOMB EFFECTS (U)

H. L. Brode

RM-1583-1

ASTIA Document Number AD 144275

Revised 3 February 1956

Copy No. 192

This material contains information affecting the national defense of the United States within the meaning of the espionage laws, Title 18 U.S.C., Secs. 793 and 794, the transmission or the revelation of which in any manner to an unauthorized person is prohibited by law.

Assigned to _____

This is a working paper. Because it may be expanded, modified, or withdrawn at any time, permission to quote or reproduce must be obtained from RAND. The views, conclusions, and recommendations expressed herein do not necessarily reflect the official views or policies of the United States Air Force.

DISTRIBUTION RESTRICTIONS

The **RAND** *Corporation*

1700 MAIN ST. • SANTA MONICA • CALIFORNIA

SECRET

57AA

4665

CLOSE-IN H-BOMB EFFECTS

I. INTRODUCTION

Recent concern with the chances of survival for relatively invulnerable structures and with the design of protective structures has led us to consider the damage due to blast, thermal, and nuclear radiations from an H-bomb in regions of greater intensity than have been previously considered important to military effects. Some aspects of the nuclear explosion remain unchanged in quality and merely increase in intensity as one moves closer to the bomb burst. Other phenomena, however, change their character sufficiently to warrant a special re-evaluation.

For example the nature of nuclear radiations may be expected to be somewhat different. The mean energy, and so the penetrating power and biological effectiveness of neutrons and gamma rays will be increased, and at the same time the rate and direction of dose delivery will be altered. Also in this close-in region the type of weapon and the details of its nucleonics will be of even greater concern. In addition we may expect large hydrodynamic effects on these doses.

In these close-in regions one may expect quite different thermal radiations. Not only will the frequency and time distributions of radiative flux be quite different than that at large distances but here any point will also be engulfed in the superheated air of the fireball itself.

SECRET

RM-1583-1

2-3-56

-2-

FORMERLY RESTRICTED DATA

Handle as Restricted Data in foreign dissemination
Section 1444b, Atomic Energy Act, 1954

Here, too, the nature of the blast wave is changed. Such quantities as positive phase duration or total impulses cannot enjoy the same significance that they do at larger distances. Aside from complications due to interaction with radiative processes and the break-down of the ideal gas assumption for air, we encounter radically different time and space histories of the blast parameters. Pressure pulses are exceedingly rapid in both rise and decay and have much greater dynamic effects. Here also we encounter hypersonic flow problems. On the other hand non-ideal surface effects may be reduced. (Strong shocks tend to be less influenced by local irregularities.) It is conceivable that precursor phenomena may not have as large an influence on the blast waves. Although the interactions between the various weapon effects is strongest in this close-in region, and although the details of bomb physics and bomb design are of greater significance here, the general nature of the free-field blast wave may still be adequately well described by a simple hydrodynamic model (using initial conditions compatible with the earlier radiative effects). The next section contains the results of detailed computations on such a model.

II. BLAST

At present measurements in the region of high levels of pressure are scanty and restricted to indirect or simple peak value measures such as fireball growth photography, time-of-arrival data and ball-crusher or beer-can type gauges. The very nature of this region has heretofore precluded extensive measurements and consequently no details of dynamic effects and time histories of pressure pulses are available from tests. For this reason, the curves offered in this section cannot be claimed to be verified in detail.

SECRET

FORMERLY RESTRICTED DATA

SECRET

RM-1583-1

2-3-56

-3-

Handle as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

However, whenever comparisons with test results are available their satisfactory agreement with these theoretical solutions speaks well for the reliability of both the theory and the data.

However, the important question as to how well the results of these calculations predict blast loading from a nuclear bomb explosion cannot be answered with any certainty at present. The following qualitative evaluation may be useful, however:

- 1.) In general a strong shock is less influenced by thermal gradients or variable conditions in the medium through which it propagates. For this reason one could expect sharper rises and cleaner shocks in this close-in region.
- 2.) Since strong shocks are characterized by very sharp rise and decay in both time and space the influence of multiple shocks and disturbances in the after-flow, due to local interferences and reflections, is minimized in problems of structural response.
- 3.) On the other hand, in the close-in regions the blast has not always developed far enough from its earliest stages to have washed out the influences of multiple bomb shock and initial radiation transport effects. Neglect of this is in general not serious, but in very special instances may mean overlooking a sudden increase in the main shock strength by as much as 20 per cent.
- 4.) Surface phenomena such as the thermal precursor should be of minor influence for any height of burst at these high levels of blast, since here the blast arrives before the bulk of the thermal radiation is received.

SECRET

5.) The hypersonic flow at shock and particle speeds of many Mach may also lead to peculiar loading problems, locally.

The curves presented are the result of numerical integration of the partial differential equations of gas motion. The calculational detail includes the use of a careful fit to the equation of state for air and an entirely general treatment of the gas dynamics (no restricting similarity assumptions, for instance).

In particular the curves presented here stem from a calculation whose initial conditions characterize a point source. Calculations also have been carried out which start from an isothermal sphere approximating the effect of radiation diffusion on the early stages of a nuclear bomb explosion. The essential character of the subsequent blast wave in these latter calculations matches that of the point-source calculation. In detail, however, the isothermal-sphere calculations are encumbered by the presence of multiple minor shocks and rarefactions. For this reason it was felt that the point-source case was more appropriate from the standpoint of simplicity and generality and for ease in scaling and analytical fitting.

The results of these calculations which include the equation of state of air do not differ from the previous ideal-gas calculations in the lower pressure regions ($\Delta P_g < 3$ atmos) but they show some marked differences in dynamic pressures, densities and temperatures in the higher pressure regions.

The general methods employed in these calculations are similar to those described in an article by this author which appeared in the Journal of Applied Physics⁽¹⁾ (June 1955) and which covers the ideal-gas computations.

All the curves are scaled to a two megaton free-air burst, with the intent that they should be used as representative of a one megaton surface burst.

¹ H. L. Brode, J. of Appl. Phys. 26 766-775 (June 1955).

SECRET

RM-1583-1
2-3-56
-5-

They also represent burst in a homogeneous sea-level atmosphere with standard ambient pressure $P_0 = 14.7$ psi, density $\rho_0 = .00129$ gm/cc, and temperature $T_0 = 273.2$ °Kelvin.

Any other yield or ambient conditions are achieved by multiplying all distances by $(W/\pi)^{1/3}$ and all times by $(W/\pi)^{1/3} \tau^{-1/2}$ where W is the yield in megatons, π is the ratio of ambient pressure (P_a) to standard pressure (P_0), and τ is the ratio of ambient temperature (T_a) to standard temperature (T_0). Under this scaling the various pressure ratios will be the same, but pressures must be multiplied by π , velocities (Mach numbers) by $\tau^{1/2}$ and temperatures by τ^1 .

Figure 1. is a plot of peak overpressure in atmospheres (14.7 psi) versus range in kilofeet.

Figure 2. is a similar plot of the peak dynamic pressure ($1/2 \rho u^2$) in atmospheres versus range in kilofeet. It should be noted that the usual Hugoniot relation for an ideal gas of $\gamma = 1.4$, viz. $Q_s = \frac{5}{2} \frac{\Delta P_s^2}{7P_0 + \Delta P_s}$ is valid only for small pressures ($\Delta P_s < 3$ atmos), and that for higher pressures the peak dynamic pressures are greater than this (characteristic of a smaller γ).

Figure 3. is a plot of particle velocity (u_s) and shock velocity (U_s) in units of ambient sonic velocity, i.e., a plot of particle and shock Mach numbers. Also plotted is the shock density (ρ_s) in units of ambient density (ρ_0) versus range in kilofeet.

Figures 4 - 21 are plots of overpressure (ΔP) in atmos, dynamic pressure (Q) in atmos, particle Mach number (U), density ratio (ρ), and temperature ratio

SECRET

FORMERLY RESTRICTED DATA

and is Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

SECRET

RM-1583-1

2-3-56

-6-

$(\theta = T/T_0)$ versus time in seconds at various distances from the burst.

Figure 22. is a plot of various parameters versus the shock overpressure (ΔP_s) in atmos. Definition of these parameters is as follows:

Q_s = shock dynamic pressure $(1/2 \rho_s U_s^2)$

t_s = time of arrival of shock (sec)

D_P^+ = duration of pressure positive phase (sec)

D_Q^+ = duration of dynamic-pressure positive phase (sec)

Δ_P^s = duration of the right-triangular pulse having the same initial slope as the true pressure pulse (sec)

Δ_Q^s = duration of the right-triangular dynamic-pressure pulse having the same initial slope as the true dynamic pressure pulse (sec)

α = exponent coefficient for early part of pressure pulse fitted by the form $\Delta P = \Delta P_s e^{-\alpha t}$

β = exponent coefficient for early part of dynamic-pressure pulse fitted by the form $Q = Q_s e^{-\beta t}$

Figures 23 - 26 show some of the blast parameters as a function of radius for various times in the course of the explosion.

SECRET

FORMERLY RESTRICTED

11A


SECRET

Handle as Restricted Data in foreign dissemination.
Section 144b, Atomic Energy Act, 1954

RM-1583-1

2-3-56

-7-



SECRET

FORMERLY RESTRICTED DATA

SECRET


FM-1583-1

2-3-56

-8-

Handle as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954


NAVY 100- KUGEL & GUNTER CO.
Incorporated 2-2-56



SECRET

SECRET

RM-1583-1
 2-3-56
 -9-



SECRET

RM-1583-1
2-3-56
-10-

as Restricted Data as being disseminated on
March 14th, Atomic Energy Act, 1954

Blast Parameters as Functions of Time at 695 Feet from a 1 MT Surface Burst


$$Q_g = 4.39 \times 10^3 \text{ atm-cs.}$$

$$t_d = .00807 \text{ sec.}$$

$$T_g = 9, T_c = 1.37 \times 10^4 \text{ K}$$

07-8-40

$$v_s = v_{s0} = 2.97 \times 10^4 \text{ feet/sec.}$$



SECRET


 4 CASSELL RD DIVISION
 KENNESAW, OHIO 44240
 NEWTON, MASS 02459
 NEWTON, MASS 02459

18-0336
SMITHKLINE BEECHAM
CONSUMER & RETAIL
PHARMACEUTICALS
1000 MARKET STREET
PHILADELPHIA, PA 19104

SECRET

RM-1583-1

2-3-56

12


Figure 6

~~Blast Parameters as Functions of Time~~
~~at 874 feet from a 1 MT Surface Burst~~

AP_s = 452 atmcs.

$$P_3 = 2.47 \times 10^3 \text{ atmos.}$$
$$t_g = .0114 \text{ sec.}$$
$$T_g = \theta_s T_0 = 7.79 \times 10^3 \text{ } ^\circ\text{K}$$

p. 10.1

$$v_s = u_{s0} = 2.03 \times 10^4 \text{ feet/sec.}$$


SECRET

FORMERLY RESTRICTED DATA

SECRET

RM-1583-1

2-3-56

-13-

Figure 7

Blast Parameters as Functions of Time
at 994 feet from a 1 MT Surface Burst

$$\Delta P_s = 297 \text{ atmos.}$$


$$Q_s = 1.55 \times 10^3 \text{ atmos.}$$

$$t_s = .0200 \text{ sec.}$$

$$T_s = \theta_s T_0 = 6.19 \times 10^3 \text{ } ^\circ\text{K}$$

$$P_s = 9.67$$

$$v_s = u_s c_s = 1.64 \times 10^4 \text{ feet/sec.}$$



SECRET

1800 - CINCINNATI
KATIE L. P. EPPER CO.
REPRODUCTION
XEROX & CO. DIVISION

FORMERLY RESTRICTED DATA


SECRET

RM-1583-1
2-3-56

as Restricted Data in Atomic Energy Act, 1954
Section 144b, Atomic Energy Act, 1954

Figure 8

Blast Parameters as Functions of Time
at 1120 feet from a 1 Mt Surface Burst



SECRET

FORMERLY RESTRICTED DATA

SECRET

RM-1583-1

2-3-56

-15-

10³ No Restricted Data in Foreign Dissemination
Section 1442, Atomic Energy Act, 1954

Figure 9

Blast Parameters as Functions of Time
at 1395 feet from a 1 MT Surface Burst

$$\Delta P_s = 99.9 \text{ atmos.}$$


$$Q_s = 385 \text{ atmos.}$$

$$t_s = .0179 \text{ sec.}$$

$$T_s = 0.5 T_0 = 3.61 \times 10^3 \text{ } ^\circ\text{K}$$

$$\rho_s = 7.36$$

$$v_s = u_s c_0 = 9.40 \times 10^3 \text{ feet/sec.}$$



MADE IN U.S.A.
21P Lines Reducing
2000 Poles, 10 to the 1000
2001 SIC KENNER & LORR CO.

SECRET

FORMERLY RESTRICTED DATA

SECRET

RM-1583-1
2-3-56

As Restricted Data in foreign disc
Section 148b, Atomic Energy Act, 1954

Figure 10

Blast Parameters as Functions of Time
at 1621 feet from 1 MB Surface Burst

$$\Delta P_s = 69.9 \text{ atmos.}$$


$$Q_s = 527 \text{ atmos.}$$

$$t_s = .0699 \text{ sec.}$$

$$T_s = 8T_0 = 2.66 \times 10^3 \text{ }^\circ\text{K}$$

$$P_s = 6.36$$

$$v_s = u_s c_0 = 7.47 \times 10^3 \text{ feet/sec.}$$




SECRET

SECRET

FORMERLY RESTRICTED DATA

Handle as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954



SECRET

Time (sec.)

0.02 0.04 0.06 0.08 0.10 0.12

0 20 40 60 80

0 20 40 60 80

SECRET

RM-1583-1

2-3-56

-18-

Figure 12

FORMERLY RESTRICTED DATA

Ends as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

Blast Parameters as Functions of Time
at 2178 feet from a 1 MT Surface Burst

$$AP_s = 30.6 \text{ atmos.}$$

$$u_s = 72.8 \text{ atmos.}$$


$$t_s = .114 \text{ sec.}$$

$$T_s = \theta_s T_0 = 1.56 \times 10^3 \theta_s$$

$$\rho_s = 5.12$$

$$v_s = u_s \theta_s = 4.76 \times 10^3 \text{ feet/sec.}$$

θ
 AP
 Q



SECRET

Time (sec.)

FORMERLY RESTRICTED DATA

Handle as Restricted Data in foreign dissemination
Section 1.44b, Atomic Energy Act, 1954

Figure 13

Blast Parameters as Functions of Time
at 2508 feet from a 3 MT Surface Burst

$$\Delta P_s = 20.7 \text{ atmos.}$$


$$v_s = 41.4 \text{ atmos.}$$

$$t_s = .205 \text{ sec.}$$

$$T_s = 1.19 \times 10^3 \text{ } ^\circ\text{K} = \theta_s T_o$$

$$P_s = 4.91$$

$$v_s = u_s c_o = 3.77 \times 10^3 \text{ feet/sec.}$$




SECRET

FORMERLY RESTRICTED DATA

Handle as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

OFFICE HEADQUARTERS
SECRET




RM-1583-1
2-3-56

P O U

SECRET

Figure 15

Blast Parameters as Functions of Time
at 3183 feet from a 1 MT Surface Burst



RM-1583-1
2-3-56
-21-0

SECRET

FORMERLY RESTRICTED DATA **SECRET**

RM-1583-1
2-3-56

As Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

Figure 16

Blast Parameters as Functions of Time
at 3636 feet from a 1 MT Surface Burst

$\Delta P_s = 7.34 \text{ atmos.}$

$Q_s = 9.46 \text{ atmos.}$

$t_s = .494 \text{ sec.}$

$T_s = \theta_s T_0 = 644^\circ \text{K}$

$\rho_s = 3.58$

$v_s = u_{sc} = 2.13 \times 10^3 \text{ feet/sec.}$





Figure 17

Blast Parameters as Functions of Time
at 441 feet from a 1 MT Surface Burst




RM-1583-1
2-3-56
23

SECRET

SECRET

FORMERLY RESTRICTED DATA

Note as Restricted Data in foreign dissemination




RM-1583-1
2-3-56
-24-3

SECRET

FORMERLY RESTRICTED DATA

as Restricted Data in foreign dissemination
Section 1440, Atomic Energy Act, 1954
Figure 19

Blast Parameters as Functions of Time
at 6197 feet from a 1 MT Surface Burst



FORMERLY RESTRICTED DATA

SECRET

24-1583-1

2-3-56

-26-

Handle as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

Figure 20

Blast Parameters as Functions of Time
at 8704 feet from a 1 Mt Surface Burst

$$AP_s = .967 \text{ atmos.}$$


$$Q_s = .314 \text{ atmos.}$$

$$L_s = 3.05 \text{ sec.}$$

$$T_s = 65T_0 = 336^\circ K$$

$$\rho_s = 1.61$$

$$v_s = u_{s0} = 573 \text{ feet/sec.}$$



Time (sec.)

SECRET

DATE 10-1-56 BY 304
RECEIVED 10-1-56
ATTACHMENT 20

SECRET

FORMERLY RESTRICTED DATA
 Handle as Restricted Data in foreign dissemination
 Section 144b, Atomic Energy Act, 1954




Figure 21

Blast Parameters as Functions of Time
 at 1589 feet from a 1 MT Surface Burst

- $\Delta P_s = 0.497$ atmos.
- $Q_s = 0.0898$ atmos.
- $t_s = 5.68$ sec.
- $T_s = \theta_s = 30^\circ K$
- $\rho_s = 1.33$
- $v_s = u_{s0} = 338$ feet/sec.

PM-1583-1
 E-3-56
 67

SECRET

SECRET


FORMERLY RESTRICTED DATA

TABLE 10. Estimated ΔP_s vs. ΔP_0 (atmos)
Boltz 1948, revised 1950, Apr. 1950

Figure 22

Various Parameters versus ΔP_0 (atmos)

(see text, page 5)



SECRET

Peak Overpressure (ΔP_0) (atmos)

SECRET

FORMERLY RESTRICTED DATA

NSA-1073-1
1-70

Figure 21

Plot Parameters as a Function of Radius
at 0.0699 sec. from a 1 MB Surface Burst


$$\Delta P_0 = 69.9 \text{ kbars.}$$

$$Q_0 = 227. \text{ g/cm}^3$$

$$P_0 = 16.86 \text{ kbar}$$

$$G_0 = 5.79 \times \frac{2469 \text{ g}}{10}$$

$$u_s = 6.87 = 2470 \text{ feet/sec.}$$



SECRET

SECRET

FORMERLY RESTRICTED DATA

shown as Restricted Data in Strength Recommendations
Section 1440, Atomic Energy Act, 1954

SM-1583-1

2-3-50

- 32 -

Figure 16

Heat Parameters as a Function of Radius
at 0.49% str. from a 1 MT Surface Burst


$$A_2 = 7.24 \text{ oct./in.}^3$$

$$Q_2 = 9.46 \text{ oct./in.}^3$$

$$G_2 = 3.52 \text{ p.}$$

$$\Theta_2 = 2.36 = \frac{6.4 \times 3 \times 10^8}{76}$$

$$u_2 = 1.96 = \frac{2.13 \times 10^8 \text{ feet/sec.}}{10^8}$$



SECRET

SECRET

RM-1583-S

10-10-56

- 33.-

FORMERLY RESTRICTED DATA

Classified as Restricted Data in foreign dissemination
Section 144b, Atomic Energy Act, 1954

SUPPLEMENT TO RM-1583-1

CLOSE-IN H-BOMB EFFECTS

The following figures (Figure 27 and Figure 28) together with the data below represent the relations between the overpressure impulse, the dynamic impulse, the peak overpressure and the distance from a one MT surface burst. The impulses are defined as follows:

$$I_P^+ = \int_{t_s}^{D_P^+} \Delta P dt \quad (\text{atmos} - \text{sec.})$$

$$I_Q^+ = \frac{1}{2} \int_{t_s}^{D_u^+} \rho u^2 dt \quad (\text{atmos} - \text{sec.})$$

ΔP (atmos)	I_P^+	I_Q^+	R(kft)
900	7.84	2.53	.695
699	7.41	2.47	.755
452	6.55	2.27	.874
297	5.81	2.09	.994
203	5.30	1.94	1.120
99.9	4.43	1.76	1.395
69.9	3.84	1.61	1.62
52.7	3.51	1.51	1.78
30.6	2.84	1.32	2.17
20.7	2.37	1.22	2.51
14.8	2.07	1.12	2.82
10.6	1.75	1.03	3.19
7.34	1.46	0.913	3.64
4.91	1.18	0.768	4.24
3.03	0.960	0.549	5.30
2.02	0.834	0.372	6.20
0.987	0.617	0.152	8.70
0.497	0.460	0.0607	12.59


SECRET

SECRET

RM-1583-1
10-10-56


FORMERLY RESTRICTED DATA

Handle as Restricted Data in Foreign Dissem
Section 144b, Atomic Energy Act, 195



Positive Impulses as a Function of the
Distance from a 1 MT Surface Burst

FORMERLY RESTRICTED DATA



SECRET

UNCLASSIFIED

UNCLASSIFIED